

seconds duration), for which treatment with aspirin was initiated.

Neurological examination was unrevealing, apart from post-traumatic left-sided amblyopia with nonparalytic strabismus (acquired during childhood).

However, we performed a brain MRI, which highlighted an extensive left-sided extra-axial accumulation of fluid, without diffusion restriction, without a well-defined capsule, measuring 15.5/5.6 centimeters in the axial plane and 10.5 centimeters in the cranio-caudal plane, with important mass effect, partial collapse of the left ventricle and midline shift of 9 mm with subfalcine herniation (see Figure 1). The lesion was classified as a type III Galassi arachnoid cyst. Subsequent neurosurgical evaluation established that it did not represent a contraindication for liver transplantation.




FIGURE 1. Left-sided extra-axial gigantic accumulation of fluid with CSF-signal on T2, FLAIR and T1 IR, determining mass effect, midline shift and subfalcine herniation

DISCUSSIONS

The imaging diagnosis of arachnoid cysts is pretty straightforward, since the fluid collection has the same density or intensity as the CSF (depending on whether CT or MRI is used), and doesn't enhance with any contrast administration. The main differential diagnosis is with epidermoid

cysts, and this is best settled by MRI. The arachnoid cyst suppresses completely with FLAIR and shows no diffusion restriction on DWI, while the epidermoid cyst does the opposite, that is, it appears hyperintense on both FLAIR and DWI. Other diagnostic considerations are chronic subdural hematoma and subdural hygroma, (which are usually bilateral and crescent shaped) or cystic meningioma (which shows ring enhancement) [3,7].

It should be noted that arachnoid cysts are common lesions, usually found incidentally during imaging procedures, and the vast majority of them are asymptomatic.

For example, Al-Holou et al analyzed retrospectively 48417 patients who underwent brain MRI and detected arachnoid cysts in 661 of them, accounting for a prevalence of 1.4%. Of the 661 patients, only 35 were considered symptomatic (~5.3%) [8].

Arachnoid cysts can give rise to intracranial hypertension, and sometimes to focal neurological deficits [1]. Sellar and suprasellar cysts are more likely to become symptomatic, while those located in the middle cranial fossa are less likely [8]. In the pediatric patients, headache is the most frequently reported symptom [9].

Other complications include intracystic hemorrhage, or rupture of the cyst into the subdural space, leading to development of a subdural hygroma [10]. Intracystic hemorrhage can also rupture and give rise to subdural hematoma [11].

Arachnoid cysts of the middle cranial fossa (which are the most common) can be graded into three types. This classification was proposed by Galassi in 1982 and takes into account their dimensions and extent, as well as their communication with the subarachnoid space (see Table 1) [12].

TABLE 1. Galassi classification of middle cranial fossa arachnoid cysts

Type	Characteristics
I	Small and limited to the anterior portion of the middle cranial fossa Free communication with the subarachnoid space.
II	Extend along the sylvian fissure and can displace the temporal lobe Slow communication with the subarachnoid space.
III	Large, fill the whole middle cranial fossa; there is displacement of multiple lobes and often there is midline shift. Little communication with the subarachnoid space.

Usually, if the cysts are asymptomatic, they should be left alone and monitored [1]. Otherwise, different neurosurgical approaches exist for symptomatic cysts, the most common being microsurgical fenestration through craniotomy, endoscopic fenestration and cystoperitoneal shunting [4]. Endoscopic techniques have been used successfully in this regard, with resolution or improvement of symptoms in the majority of patients, and no peri-operative mortality or significant morbidity [13]. Other authors however, have noted a high proportion of relapses after endoscopic fenestration and achieved better results with microsurgical fenestration [14]. The choice of neurosurgical treatment is also influenced by the location of

arachnoid cysts. In one study, the best results were obtained by using fenestration with cystoperitoneal shunting for supratentorial cysts, endoscopic fenestration for midline cysts and cystoperitoneal shunting for infratentorial cysts [15].

CONCLUSIONS

Arachnoid cysts are usually asymptomatic lesions, found incidentally on imaging procedures. However, they can grow to enormous size, at times causing mass effect and compressing nearby structures. Periodic neurological and imaging assessment is therefore recommended.

REFERENCES

- Allan HR, Martin AS, Joshua PK, Sashank P. Intracranial Neoplasms and Paraneoplastic Disorders. In: Allan HR, Martin AS, Joshua PK, Sashank P. Adams and Victor's Principles of Neurology – 11th Edition. McGraw-Hill Educational, 2019:694.
- Westermaier T, Schweitzer T, Ernestus RI. Arachnoid cysts. *Adv Exp Med Biol.* 2012;724:37-50.
- Kress B. Cysts. In: Sartor K, Haehnel S, Kress B. Brain Imaging (Direct Diagnosis in Radiology). Thieme, 2008:181-183.
- Mustansir F, Bashir S, Darbar A. Management of Arachnoid Cysts: A Comprehensive Review. *Cureus.* 2018;10(4):e2458.
- Srinivasan US, Lawrence R. Posterior fossa arachnoid cysts in adults: Surgical strategy: Case series. *Asian J Neurosurg.* 2015;10(1):47.
- Dyck P, Gruskin P. Supratentorial arachnoid cysts in adults. A discussion of two cases from a pathophysiologic and surgical perspective. *Arch Neurol.* 1977 May;34(5):276-279.
- Osborn AG. Arachnoid Cyst. In Osborn A.G. et al. Diagnostic Imaging: Brain, First Edition. Amrystis, 2004: 1-7-4.
- Al-Holou WN, Terman S, Kilburg C, et al. Prevalence and natural history of arachnoid cysts in adults. *J Neurosurg.* 2013; 118(2):222-231.
- Mazurkiewicz-Beldzińska M, Dilling-Ostrowska E. Presentation of intracranial arachnoid cysts in children: correlation between localization and clinical symptoms. *Med Sci Monit.* 2002; 8(6):CR462-5.
- Jafrani R, Raskin JS, Kaufman A, Lam S. Intracranial arachnoid cysts: Pediatric neurosurgery update. *Surg Neurol Int.* 2019;10:15.
- Hall A, White MAJ, Myles L. Spontaneous subdural haemorrhage from an arachnoid cyst: a case report and literature review. *Br J Neurosurg.* 2017;31(5):607-610.
- Galassi E, Tognetti F, Gaist G, et al. CT scan and metrizamide CT cisternography in arachnoid cysts of the middle cranial fossa: classification and pathophysiological aspects. *Surg Neurol.* 1982;17(5):363-369.
- Karabatsou K, Hayhurst C, Buxton N, et al. Endoscopic management of arachnoid cysts: an advancing technique. *J Neurosurg.* 2007;106(6 Suppl):455-462.
- Holst AV, Danielsen PL, Juhler M. Treatment options for intracranial arachnoid cysts: a retrospective study of 69 patients. *Acta Neurochir Suppl.* 2012;114:267-270.
- Ahn JY, Chio JU, Yoon SH, et al. Treatment and Outcome of Intracranial Arachnoid Cysts. *J Korean Neurosurg Soc.* 1994;23(2):194-203.